Be certain to **know** the quizzes and . . .

1 Systems of Linear Equations

1. Solve the following systems of linear equations using row reduction.

(a)
$$\begin{cases} x_1 & -2x_2 & & & -6x_5 & = 0 \\ & x_2 & +x_3 & +6x_4 & & = 5 \\ & 2x_2 & & +6x_4 & +x_5 & = 4 \\ & x_2 & -x_3 & & +x_5 & = -1 \end{cases}$$

(b)
$$\begin{cases} 2x_1 & -2x_2 & +4x_3 & = 2 \\ & & x_3 & = 0 \\ x_1 & +x_2 & +2x_3 & = 0 \end{cases}$$

(c)
$$\begin{cases} 2x_1 & -2x_2 & +4x_3 & = 2 \\ -x_1 & -x_2 & +3x_3 & = 2 \\ x_1 & -3x_2 & +7x_3 & = 2 \end{cases}$$

2. Solve the following systems of linear equations by setting up problem as a matrix problem and by finding an inverse matrix.

(a)
$$\begin{cases} 2x_1 & -2x_2 & +4x_3 & = 2 \\ & -x_2 & +3x_3 & = 2 \\ & -3x_2 & +7x_3 & = 2 \end{cases}$$

(b)
$$\begin{cases} 2x_1 & -2y & = 2 \\ -x_1 & -3y & = 2 \end{cases}$$

2 Matrices, Determinants and Cramer's Rule

3. Let $A = \begin{bmatrix} 2 & -2 & 0 \\ -1 & 4 & 1 \\ 0 & 4 & 1 \end{bmatrix}$, $B = \begin{bmatrix} 2 & -2 & 0 & 0 \\ 2 & -2 & 1 & 2 \\ 2 & -2 & 0 & 3 \\ 2 & -2 & 5 & 4 \end{bmatrix}$, $C = \begin{bmatrix} 2 & -2 & 0 & -2 & 0 \\ 0 & -2 & 0 & -2 & 0 \\ 0 & 0 & 3 & -2 & 0 \\ 0 & 0 & 0 & -2 & 0 \\ 0 & 0 & 0 & 0 & 4 \end{bmatrix}$

and $D = \begin{bmatrix} 2 & 0 & 0 \\ 0 & 4 & 0 \\ 0 & 0 & 1 \end{bmatrix}$.

(a) Find the determinant of the matices A, B, C and D

(b) Compute D^3 and D^{-1}.

(c) Compute $C^T C$. What kind of matrix is $C^T C$?

4. Solve the following equations for X assuming any matrix has an inverse. Let A, B, C, X be nxn matrices and let $\mathbf{u}$ be an $nx1$ vector.

(a) $AX = BX - A$

(b) $AX = 2X - A$

(c) $A\mathbf{u} = 2\mathbf{u} + B$

5. Assume $\det(A) = 11$.

(a) Switch R1 and R2 in the matrix A to get matrix B, what is $\det(B)$?

(b) Replace R1 with R1 - 4R2 to get matrix B, what is $\det(B)$?

(c) Mutiply R1 by 4 to get matrix B, what is $\det(B)$?

6. Solve the following using Cramer's rules.

(a) $\begin{cases} 2x_1 & -2x_2 & +4x_3 & = 2 \\ & -x_2 & +3x_3 & = 2 \\ & -3x_2 & +7x_3 & = 2 \end{cases}$

(b) $\begin{cases} 2x_1 & -2y & = 2 \\ -x_1 & -3y & = 2 \end{cases}$

3   Vectors

7. Let $\mathbf{v} = (1, 3, 4)$ and $\mathbf{w} = (1, -1, 0)$ be vectors in $\mathbb{R}^3$. and let $P(1, 1, 1)$ and $Q(0, -4, 0)$ be two points in $\mathbb{R}^3$.

(a) Find a vector that is parallel to $\mathbf{v}$ and unit.

(b) Compute $\|2\mathbf{v} - \mathbf{w}\|$.

(c) Compute the angle between $\mathbf{v}$ and $\mathbf{w}$.

(d) Find the equation of a plane containing P and with normal vector $\mathbf{v}$.

8. Find parametric equations for the line (in $\mathbb{R}^3$) so that

(a) the line contains the point $P(0, 1, 2)$ and $Q(1, 2, 3)$.

(b) the line contains the point $P(-5, 1, 3)$ and is parallel to the vector $(1, 2, 3)$.

9. Find equation for the plane (in $\mathbb{R}^3$) so that

   (a) the plane contains the point $P(2, 2, 2)$, $Q(1, 2, 3)$ and $R(1, -1, 0)$.

   (b) the plane contains the origin and is perpendicular to the vector $(1, 2, 3)$.

10. Define the planes P_1 and P_2 as follows:

$$P_1 : x - 2y + z = 12$$

$$P_2 : 3x - 3y + z = 4$$

   (a) What are the two normal vectors for the above planes.

   (b) Find the angle between the two above planes.

   (c) Find two pointsone on each of the above planes.

   (d) Find set of all points that lay in both planes.